

THE STEREOCHEMISTRY OF VINYLACETYL-CoA-ISOMERASE OF *CLOSTRIDIUM KLUYVERI*

H. HASHIMOTO, H. GÜNTHER and H. SIMON

Lehrstuhl für Organische Chemie und Biochemie des Organisch-Chemischen Laboratoriums der T.U. München, D 8 München 2, Arcisstr. 21, Germany

Received 13 April 1973

1. Introduction

Vinylacetyl-CoA isomerases [EC 5.3.3.3], catalyzing the reaction **1a** → **2a** have been found in mammals [1] and microorganisms, such as *Clostridi* [2]. Little about the mechanism, and nothing about the stereochemistry of this 1,3-proton shift is known. An internal hydrogen transfer appears to be involved because tritium was not incorporated into the acyl portion of β -methyl-crotonyl-CoA when β -methylvinyl-acetyl-CoA was isomerized in H₂O/HTO by the enzyme from ox liver [1]. From the inhibitory action of *p*-chloromercuribenzoate Bartsch and Barker [2] inferred a functional thiol group. In the isomerization **1** → **2** the carbon atoms 2 and 4 are involved. Whether the pro R or pro S hydrogen atom at C-2 is eliminated and from which side the hydrogen atom at carbon atom 4 enters, was here determined by the fermentation of (R,S)- α -methylvinylacetate (**1b**) and tiglate (**2b**) with whole cells in H₂O or with Δ^3 -pentenate (**1c**) in D₂O, respectively.

2. Materials and methods

C. kluyveri cultures were originally revived from dried crotonate cells (C.F. Boehringer u. Soehne, Biochemica, Tutzing) and then grown as described [3] without changing the crotonate medium to ethanol acetate. α -Methylvinyl-acetate [4] and Δ^3 -pentenate [5] were synthesized. The purity was checked by GLC and NMR-spectroscopy.

For GLC aliquots were acidified with 0.2 N HCl and the organic acids separated on a column

(69 in. × 1/12 in.) filled with 20% neopentylglycol succinate and 2% phosphoric acid on Chromosorb P.

Compounds **1b** and **3a** were separated by counter-current distribution with *n*-hexane and water.

Other preparative separations of acids were carried out by the method of Marvel and Rands [6].

For polarometric measurements a Jasco Model ORD/UV-5 (Japan Spectroscopic) was used.

3. Results and discussion

In the presence of crotonate or butyrate, *C. kluyveri* converted (R,S)- α -methylvinylacetate (R,S **1b**) only partially to α -methylbutyrate (**3a**) (table 1). The remaining α -methylvinylacetate was isolated and was found to be the (R)(-)-enantiomer (R, **1b**) with a specific optical rotation (table 1) nearly identical to the highest reported value [7]. Obviously, reaction occurred only with the (S)(+)-enantiomer of **1b**, and its hydrogen atom (R¹ = H) had migrated. Therefore one has to assume that the pro S hydrogen atom vinylacetyl-CoA migrates to form crotonyl-CoA.

The fermentation product, [S] (+)- α -methylbutyrate (**3a**), had the optical rotation (table 1) reported before [8], in accordance with the absolute stereochemistry of the butyryl-CoA dehydrogenase recently determined by us [3]. Then it was assumed that **2b** was the intermediate. Substrate **2b** led also to **3a**, with the same specific rotation (table 1).

In the fermentation of Δ^3 -pentenate (**1c**) in D₂O, **1c** is hydrogen donor and acceptor. Obviously, one equivalent of **1c** was hydrogenated to **3b** and another was degraded via **1c** → **2c** → **4** → **5** → **6** + **7**.

Table 1
Fermentation^a of (R,S) α -methylvinylacetate (R,S1b) 2-methylcrotonate (2b) in H₂O and Δ^3 -pentenate (1c) in D₂O.


Substrate and cosubstrate (mMol)	<i>C. kluyveri</i> (g wet packed cells/ mMol substrate)	Products ^b (mMol)	Specific optical rotation
(R,S) α -methylvinylacetate (20) and butyrate (10)	1.5	(S) α -Methylbutyrate (9.6) and (R) α -methylvinylacetate (7.0)	$[\alpha]^D = +21.0^\circ$ $[\alpha]^D = -39.7^\circ$
2-Methylcrotonate (tiglate) (1.0) and crotonate (1.0)	0.75	(S) α -Methylbutyrate (0.54)	$[\alpha]^D = +21.0^\circ$
Δ^3 -Pentenate (5.0)	1.5	(R) [2- ² H]propionate ^c (1.33)	$[\alpha]^{365} = -2.0^\circ$
Δ^3 -Pentenate (2.0)	3.6	(R) [2- ² H]propionate ^d (0.42)	$[\alpha]^{365} = -1.3^\circ$

^a The cells were harvested by centrifugation (12 000 g), washed twice with 0.1 M phosphate buffer pH 7.0 and incubated (40 hr) with substrates at 37°C. All manipulations were done under nitrogen.

^b Only products of interest from a stereochemical point of view are given.

^c 0.64 deuterium atoms/mole.

^d 0.50 deuterium atoms/mole.



Scheme 1

For this reason, crotonate or butyrate was not added during the incubations. The table shows the deuterium amounts and optical rotations of propionate **6**, isolated from the experiment in D₂O. According to the determination of the absolute stereochemistry [9, 10] it had the *R* configuration. The conversion **2c** → **6** should not change the configuration of the original carbon atom 4 of **2c**. Consequently, the hydrogen atom entered from the re-side. The deuterium content of the medium was 90% D₂O, but only 0.6 deuterium atoms were incorporated. Thus, the 1.3 proton shift appears to be only partially intramolecular, in contrast to the results of Rilling and Coon [1].

Acknowledgements

It is gratefully acknowledged that Deutsche Forschungsgemeinschaft supported this work, also by a grant to Dr. H.H. We thank Miss Klüpfel for skilled technical assistance.

References

- [1] H.C. Rilling and M.J. Coon, J. Biol. Chem. 235 (1960) 3087.
- [2] R.G. Bartsch and H.A. Barker, Arch. Biochem. Biophys. 92 (1961) 122.
- [3] H.J. LaRoche, M. Kellner, H. Gunther and H. Simon, Hoppe-Seyler's Z. Physiol. Chem. 352 (1971) 399.
- [4] J.F. Lane, J.D. Roberts and W.G. Young, J. Am. Chem. Soc. 66 (1944) 545.
- [5] R.P. Linstead, E.G. Noble and E.J. Boorman, J. Chem. Soc. (1933) 557.
- [6] C.S. Marvel and R.D. Rands, Jr., J. Am. Chem. Soc. 72 (1950) 2642.
- [7] R. Rossi and G. Ingrosso, Gazz. Chim. Ital. 98 (1968) 866.
- [8] L. Lardicci and L. Conti, Ann. Chim. (Rome) 51 (1967) 823; K. Freudenberg and W. Lwowski, Liebigs Ann. Chem. 594 (1955) 76.
- [9] B. Zagalak, P.A. Frey, G.L. Karabatsos and R.H. Abeles, J. Biol. Chem. 241 (1966) 3028.
- [10] A. Retey, A. Umani-Ronchi and D. Arigoni, Experientia [Basel] 22 (1966) 72.